

Unit - 2 : Inverse Laplace Transform

Formulae :-

$$1. \quad L^{-1}\left(\frac{1}{s}\right) = 1$$

$$9. \quad L^{-1}\left[\frac{1}{s^2 - a^2}\right] = \frac{1}{a} \sinh at$$

$$2. \quad L^{-1}\left(\frac{k}{s}\right) = k$$

$$10. \quad L^{-1}\left[\frac{1}{s^{n+1}}\right] = \frac{t^n}{n!}$$

$$3. \quad L^{-1}\left(\frac{1}{s+a}\right) = e^{-at}$$

$$4. \quad L^{-1}\left(\frac{1}{s-a}\right) = e^{at}$$

$$5. \quad L^{-1}\left(\frac{a}{a^2 + s^2}\right) = \sin at$$

$$6. \quad L^{-1}\left[\frac{1}{s^2 + a^2}\right] = \frac{1}{a} \sin at$$

$$7. \quad L^{-1}\left[\frac{s}{s^2 + a^2}\right] = \cos at$$

$$8. \quad L^{-1}\left[\frac{s}{s^2 - a^2}\right] = \cosh at$$

First Shifting property :-

$$\text{IF } \mathcal{L}^{-1} [\bar{F}(s)] = F(t)$$

$$\mathcal{L}^{-1} [\bar{F}(s-a)] = e^{at} F(t) \\ = e^{at} \mathcal{L}^{-1} [\bar{F}(s)]$$

$$\mathcal{L}^{-1} [\bar{F}(s+a)] = e^{-at} F(t) \\ = e^{-at} \mathcal{L}^{-1} [\bar{F}(s)]$$

* Examples :

1. Find $\mathcal{L}^{-1} \left[\frac{4s+15}{16s^2-25} \right]$

→ Consider

$$\frac{4s+15}{16s^2-25} = \frac{4s+15}{16 \left[s^2 - \frac{25}{16} \right]}$$

$$= \frac{4s+15}{16 \left[s^2 - \left(\frac{5}{4} \right)^2 \right]}$$

$$\mathcal{L}^{-1} \left[\frac{4s+15}{16s^2-25} \right] = \frac{1}{16} \mathcal{L}^{-1} \left[\frac{4s}{s^2 - (5/4)^2} \right] + \frac{15}{16} \mathcal{L}^{-1} \left[\frac{1}{s^2 - (5/4)^2} \right]$$

$$= \frac{4}{16} \mathcal{L}^{-1} \left[\frac{s}{s^2 - (5/4)^2} \right] + \frac{15}{16} \mathcal{L}^{-1} \left[\frac{1}{s^2 - (5/4)^2} \right]$$

$$= \frac{1}{4} \cosh \frac{s}{4} + \frac{15}{16} \times \frac{1}{5/4} \mathcal{L}^{-1} \left[\frac{5/4}{s^2 - (5/4)^2} \right]$$

$$= \frac{1}{4} \cosh \frac{5}{4} t + \frac{15}{16} \times \frac{4}{5} \sinh \frac{5}{4} t$$

$$= \frac{1}{4} \cosh \frac{5}{4} t + \frac{3}{4} \sinh \frac{5}{4} t$$

2. $\mathcal{L}^{-1} \left[\frac{s+1}{s^2+s+1} \right]$

→

Consider $\frac{s+1}{s^2+s+1}$ — (I)

$$ax^2 + bx + c$$

$$a=1, b=1, c=1$$

$$\text{Complete square term} = \left(\frac{-b}{2a} \right)^2$$

$$= \left(\frac{-1}{2 \times 1} \right)^2 = \left(\frac{-1}{2} \right)^2 = \frac{1}{4}$$

eqⁿ (I) becomes

$$\frac{s+1}{s^2+s+1} = \frac{s+1}{s^2+s+\frac{1}{4}+1-\frac{1}{4}}$$

$$= \frac{s+1}{(s+\frac{1}{2})^2 + \frac{4-1}{4}}$$

$$= \frac{s+1}{(s+\frac{1}{2})^2 + \frac{3}{4}}$$

$$\frac{s+1}{s^2+s+1} = \frac{s+1}{(s+\frac{1}{2})^2 + (\frac{\sqrt{3}}{2})^2}$$

Taking L^{-1} on both

$$L^{-1} \left[\frac{s+1}{s^2+s+1} \right] = L^{-1} \left[\frac{s+1}{(s+\frac{1}{2})^2 + (\frac{\sqrt{3}}{2})^2} \right]$$

$$= L^{-1} \left[\frac{s+\frac{1}{2}+\frac{1}{2}}{(s+\frac{1}{2})^2 + (\frac{\sqrt{3}}{2})^2} \right]$$

$$L^{-1} \left[\frac{s+1}{s^2+s+1} \right] = L^{-1} \left[\frac{s+\frac{1}{2}}{(s+\frac{1}{2})^2 + (\frac{\sqrt{3}}{2})^2} \right] + \frac{1}{2} L^{-1} \left[\frac{1}{(s+\frac{1}{2})^2 + (\frac{\sqrt{3}}{2})^2} \right]$$

$$= e^{-\frac{1}{2}t} L^{-1} \left[\frac{s}{s^2 + (\frac{\sqrt{3}}{2})^2} \right] + \frac{1}{2} e^{-\frac{1}{2}t} L^{-1} \left[\frac{1}{s^2 + (\frac{\sqrt{3}}{2})^2} \right]$$

$$= e^{-\frac{1}{2}t} \cos \frac{\sqrt{3}}{2} t + \frac{1}{2} e^{-\frac{1}{2}t} \frac{1}{\frac{\sqrt{3}}{2}} \sin \frac{\sqrt{3}}{2} t$$

$$= e^{-\frac{1}{2}t} \cos \frac{\sqrt{3}}{2} t + \frac{1}{\sqrt{3}} e^{-\frac{1}{2}t} \sin \frac{\sqrt{3}}{2} t$$

$$= e^{-\frac{1}{2}t} \left[\cos \frac{\sqrt{3}}{2} t + \frac{1}{\sqrt{3}} \sin \frac{\sqrt{3}}{2} t \right]$$

3. $L^{-1} \left[\frac{15}{s^2+4s+13} \right]$

→ Consider,

$$\frac{15}{s^2 + 4s + 13} \quad \text{--- (I)}$$

$$ax^2 + bx + c$$

$$a=1, b=4, c=13$$

$$\text{Complete square term} = \left(\frac{-b}{2a} \right)^2$$

$$= \left(\frac{-4}{2 \times 1} \right)^2 = (-2)^2 = 4$$

eqⁿ (I) becomes

$$\frac{15}{s^2 + 4s + 13} = \frac{15}{s^2 + 4s + 4 + 13 - 4}$$

$$= \frac{15}{(s+2)^2 + 3^2}$$

$$\frac{15}{s^2 + 4s + 13} = \frac{15}{(s+2)^2 + 3^2}$$

Taking L^{-1} on both side

$$L^{-1} \left[\frac{15}{s^2 + 4s + 13} \right] = L^{-1} \left[\frac{15}{(s+2)^2 + 3^2} \right]$$

$$= 15 L^{-1} \left[\frac{1}{(s+2)^2 + 3^2} \right]$$

$$= 15 e^{-2t} L^{-1} \left[\frac{1}{s^2 + 9} \right]$$

$$= 15 e^{-2t} \frac{1}{3} \sin 3t$$

$$\boxed{\mathcal{L}^{-1} \left[\frac{15}{s^2+4s+13} \right] = 5 e^{-2t} \sin 3t}$$

4. $\mathcal{L}^{-1} \left[\frac{s+8}{s^2+4s+5} \right]$

→ Consider,

$$\frac{s+8}{s^2+4s+5} \quad \text{--- (1)}$$

$$ax^2 + bx + c$$

$$a=1, \quad b=4, \quad c=5$$

$$\text{Complete Square term} = \left(\frac{-b}{2a} \right)^2$$

$$= \left(\frac{-4}{2 \times 1} \right)^2 = (-2)^2 = 4$$

eqⁿ (1) becomes

$$\frac{s+8}{s^2+4s+5} = \frac{s+8}{s^2+4s+4+5-4}$$

$$= \frac{s+8}{(s+2)^2+1}$$

$$\frac{s+8}{s^2+4s+5} = \frac{s+8}{(s+2)^2+1}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{s+8}{s^2+4s+5} \right] = L^{-1} \left[\frac{s+8}{(s+2)^2+1} \right]$$

$$= L^{-1} \left[\frac{(s+2)+6}{(s+2)^2+1} \right]$$

$$L^{-1} \left[\frac{s+8}{s^2+4s+5} \right] = L^{-1} \left[\frac{s+2}{(s+2)^2+1} \right] + 6 L^{-1} \left[\frac{1}{(s+2)^2+1} \right]$$

$$= e^{-2t} L^{-1} \left[\frac{s}{s^2+1} \right] + 6e^{-2t} L^{-1} \left[\frac{1}{s^2+1} \right]$$

$$= e^{-2t} \cos t + 6e^{-2t} \sin t$$

$$= e^{-2t} \cos t + 6e^{-2t} \sin t$$

$$= e^{-2t} [\cos t + 6 \sin t]$$

~~Imp~~
*

Inverse Laplace Transform of derivative of $\bar{F}(s)$

$$L^{-1} \left[\frac{d}{ds} [\bar{F}(s)] \right] = -t \bar{F}(t)$$

$$\mathcal{L}^{-1} \left[\frac{d^2}{ds^2} [\bar{F}(s)] \right] = (-1)^2 t^2 F(t)$$

Examples

1 Find $\mathcal{L}^{-1} \left[\log \left(1 + \frac{1}{s^2} \right) \right]$

Consider

$$\bar{F}(s) = \log \left[1 + \frac{1}{s^2} \right]$$

$$= \log \left[\frac{s^2+1}{s^2} \right]$$

$$= \log [s^2+1] - \log [s^2]$$

diff. w.r.t. 's'

$$\frac{d}{ds} [\bar{F}(s)] = \frac{d}{ds} [\log (s^2+1) - \log (s^2)]$$

$$= \frac{1}{s^2+1} \frac{d}{ds} (s^2+1) - \frac{1}{s^2} \frac{d}{ds} (s^2)$$

$$= \frac{2s}{s^2+1} - \frac{2s}{s^2}$$

$$\frac{d}{ds} [\bar{F}(s)] = \frac{2s}{s^2+1} - \frac{2}{s}$$

taking L^{-1} on both

$$L^{-1} \left[\frac{d}{ds} [\bar{F}(s)] \right] = L^{-1} \left[\frac{2s}{s^2+1} - \frac{2}{s} \right]$$

$$-tF(t) = 2L^{-1} \left[\frac{s}{s^2+1} \right] - 2L^{-1} \left[\frac{1}{s} \right]$$

$$-tF(t) = 2 \cos t - 2 \times 1$$

$$tF(t) = 2 - 2 \cos t$$

$$F(t) = \frac{2(1 - \cos t)}{t}$$

2. Find $L^{-1} \left[\log \left(\frac{s+a}{s+b} \right) \right]$

→ Solⁿ, let $\bar{F}(s) = \log \left(\frac{s+a}{s+b} \right)$

$$= \log(s+a) - \log(s+b)$$

DIFF. w.r.t.s

$$\frac{d}{ds} [\bar{F}(s)] = \frac{d}{ds} [\log(s+a) - \log(s+b)]$$

$$= \frac{1}{s+a} \times 1 - \frac{1}{s+b} \times 1$$

$$= \frac{1}{s+a} - \frac{1}{s+b}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{d}{ds} \bar{F}(s) \right] = e^{-at} - e^{-bt}$$

$$t F(t) = e^{-at} - e^{-bt}$$

$$F(t) = \frac{e^{-at} - e^{-bt}}{t}$$

3) Find $L^{-1} \left[\log \left(\frac{s+1}{s-1} \right) \right]$

→

$$\text{Let } \bar{F}(s) = \log \left(\frac{s+1}{s-1} \right)$$

$$\bar{F}(s) = \log(s+1) - \log(s-1)$$

DIFF. w.r.t. s

$$\frac{d}{ds} [\bar{F}(s)] = \frac{d}{ds} [\log(s+1) - \log(s-1)]$$

$$t F(t) = \frac{1}{s+1} - \frac{1}{s-1}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{d}{ds} (\bar{F}(s)) \right] = L^{-1} \left[\frac{1}{s+1} - \frac{1}{s-1} \right]$$

$$tF(t) = e^{-t} - e^t$$

$$F(t) = \frac{e^t - e^{-t}}{t}$$

Imp

4. Find $L^{-1} \left[\frac{\cot^{-1} (s+3)}{2} \right]$

→ Let $\bar{F}(s) = \cot^{-1} \left(\frac{s+3}{2} \right)$

diff w.r.t s

$$\frac{d}{ds} [\bar{F}(s)] = \frac{d}{ds} \left[\cot^{-1} \left(\frac{s+3}{2} \right) \right]$$

$$\text{use } \frac{d}{dx} [\cot^{-1}(x)] = \frac{-1}{1+x^2} \frac{d}{dx} (x)$$

$$= \frac{-1}{1 + \left(\frac{s+3}{2} \right)^2} \frac{d}{ds} \left(\frac{s+3}{2} \right)$$

$$= \frac{-1}{1 + \frac{(s+3)^2}{4}} \times \frac{1}{2} (1+0)$$

$$= \frac{-1}{4 + \frac{(s+3)^2}{4}} \times \frac{1}{2}$$

$$= \frac{-4^2}{4 + (s+3)^2} \times \frac{1}{2}$$

$$= \frac{-2}{4 + (s+3)^2}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{d}{ds} [\bar{F}(s)] \right] = L^{-1} \left[\frac{-2}{4 + (s+3)^2} \right]$$

$$-tF(t) = -2 L^{-1} \left[\frac{1}{4 + (s+3)^2} \right]$$

$$-tF(t) = -2 e^{-3t} L^{-1} \left[\frac{1}{4 + s^2} \right]$$

$$-tF(t) = -2 e^{-3t} \frac{1}{2} \sin 2t$$

$$F(t) = \frac{e^{-3t} \sin 2t}{t}$$

H.W.
5. Find $L^{-1} \left[\cot^{-1} \left(\frac{s}{2} \right) \right]$

→ Let $\bar{F}(s) = \cot^{-1} \left(\frac{s}{2} \right)$

diff. w.r.t. s

$$\frac{d}{ds} [\bar{F}(s)] = \frac{d}{ds} \left[\cot^{-1} \left(\frac{s}{2} \right) \right]$$

$$\text{use } \frac{d}{dx} [\cot^{-1} x] = \frac{-1}{1+x^2}$$

$$\frac{d}{ds} = \frac{-1}{1+\left(\frac{s}{2}\right)^2} \cdot \frac{d}{ds} \left(\frac{s}{2} \right)$$

$$= \frac{-1}{1+s^2/4} \cdot \frac{1}{2}$$

$$= \frac{-1}{\frac{4+s^2}{4}} \times \frac{1}{2}$$

$$= \frac{-4}{4+s^2} \times \frac{1}{2}$$

$$= \frac{-2}{4+s^2}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{d}{ds} [\bar{F}(s)] \right] = L^{-1} \left[\frac{-2}{4+s^2} \right]$$

$$-t F(t) = -2 \cdot L^{-1} \left[\frac{1}{s^2+4} \right]$$

$$-t F(t) = -2 \times \frac{1}{2} \sin 2t$$

$$t F(t) = -\sin 2t$$

$$F(t) = \frac{\sin 2t}{t}$$

6. Find $L^{-1} \left[\tan^{-1} \left(\frac{2}{s^2} \right) \right]$

→

$$\text{Let } \bar{F}(s) = \tan^{-1} \left(\frac{2}{s^2} \right)$$

DIFF. w.r.t. s.

$$\frac{d}{ds} [\bar{F}(s)] = \frac{d}{ds} \left[\tan^{-1} \left(\frac{2}{s^2} \right) \right]$$

$$= \frac{1}{1 + \left(\frac{2}{s^2} \right)^2} \frac{d}{ds} \left(\frac{2}{s^2} \right)$$

$$= \frac{1}{1 + \frac{4}{s^4}} \left(\frac{2(-2)}{s^3} \right)$$

$$= \frac{1}{\frac{s^4+4}{s^4}} \times \frac{-4}{s^3}$$

$$= \frac{s^4}{s^4+4} \times \frac{-4}{s^3}$$

$$= \frac{-4s}{s^4+4}$$

$$\frac{d}{ds} [\bar{F}(s)] = \frac{-4s}{s^4+4}$$

$$\frac{d}{ds} [\bar{F}(s)] = \frac{1}{(s+1)^2+1} - \frac{1}{(s-1)^2+1}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{d}{ds} [\bar{F}(s)] \right] = L^{-1} \left[\frac{1}{(s+1)^2+1} - \frac{1}{(s-1)^2+1} \right]$$

$$= L^{-1} \left[\frac{1}{(s+1)^2+1} \right] - L^{-1} \left[\frac{1}{(s-1)^2+1} \right]$$

$$= e^{-t} L^{-1} \left[\frac{1}{s^2+1} \right] - e^t L^{-1} \left[\frac{1}{s^2+1} \right]$$

$$\therefore t F(t) = e^{-t} \sin t - e^t \sin t$$

$$t F(t) = e^t \sin t - e^{-t} \sin t$$

$$= \sin t [e^t - e^{-t}]$$

$$t F(t) = 2 \sin t \left(\frac{e^t - e^{-t}}{2} \right)$$

$$t F(t) = 2 \sin t \cdot \sinh t$$

$$\therefore \boxed{F(t) = \frac{2 \sin t \sinh t}{t}}$$

Imp

* Partial Fraction.

$$1. \frac{ax^2+bx+c}{(x-1)(x+2)(x-3)} = \frac{A}{(x-1)} + \frac{B}{(x+2)} + \frac{C}{(x-3)}$$

$$2. \frac{ax^2+bx+c}{(x^2+2)(x+3)} = \frac{Ax+B}{(x^2+2)} + \frac{C}{(x+3)}$$

$$3. \frac{ax^2+bx+c}{(x^2+2x+1)(x+4)} = \frac{Ax+B}{(x^2+2x+1)} + \frac{C}{(x+4)}$$

$$4. \frac{ax^2+bx+c}{(x+1)^3} = \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{C}{(x+1)^3}$$

Examples :-

$$1. \text{ Find } L^{-1} \left[\frac{s^2+2s-3}{s(s-3)(s+2)} \right]$$

Consider

$$\frac{s^2+2s-3}{s(s-3)(s+2)} = \frac{A}{s} + \frac{B}{(s-3)} + \frac{C}{(s+2)} \quad \text{--- (2)}$$

$$\frac{s^2+2s-3}{s(s-3)(s+2)} = \frac{A(s-3)(s+2) + B \cdot s(s+2) + C \cdot s(s-3)}{s(s-3)(s+2)}$$

$$s^2+2s-3 = A(s-3)(s+2) + B \cdot s(s+2) + C \cdot s(s-3)$$

$$s^2+2s-3 = A(s^2+2s-3s-6) + B(s^2+2s) + C(s^2-3s)$$

$$s^2+2s-3 = A(s^2-s-6) + B(s^2+2s) + C(s^2-3s)$$

$$s^2 + 2s - 3 = As^2 - As - 6A + Bs^2 + B2s + Cs^2 - 3Cs$$

$$s^2 + 2s - 3 = s^2(A+B+C) + s(-A+2B-3C) - 6A$$

Equating powers of s

$$A + B + C = 1$$

$$-A + 2B - 3C = 2$$

$$-6A = -3$$

$$\boxed{A = \frac{1}{2}}$$

$$B + C = 1 - A$$

$$B + C = 1 - \frac{1}{2}$$

$$\boxed{B + C = \frac{1}{2}}$$

$$2B - 3C = 2 + A$$

$$= 2 + \frac{1}{2}$$

$$\boxed{2B - 3C = \frac{5}{2}}$$

$$2B - 3C = \frac{5}{2}$$

$$3B + 3C = \frac{3}{2}$$

$$5B + 0 = \frac{5}{2} + \frac{3}{2}$$

$$5B = \frac{8}{2} = 4$$

$$\therefore \boxed{B = \frac{4}{5}}$$

$$B + C = \frac{1}{2}$$

$$C = \frac{1}{2} - B$$

$$C = \frac{1}{2} - \frac{4}{5}$$

$$C = \frac{S-8}{10}$$

$$C = \frac{-3}{10}$$

eqⁿ (i) becomes

$$\frac{S^2 + 2S - 3}{S(S-3)(S+2)} = \frac{1}{2} \times \frac{1}{S} + \frac{4}{5} \times \frac{1}{S-3} + \left(\frac{-3}{10}\right) \times \frac{1}{S+2}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{S^2 + 2S - 3}{S(S-3)(S+2)} \right] = \frac{1}{2} L^{-1} \left[\frac{1}{S} \right] + \frac{4}{5} L^{-1} \left[\frac{1}{S-3} \right] - \frac{3}{10} L^{-1} \left[\frac{1}{S+2} \right]$$

$$= \frac{1}{2} \times 1 + \frac{4}{5} e^{3t} - \frac{3}{10} e^{-2t}$$

2. Find $L^{-1} \left[\frac{2S^2 - 1}{(S^2 + 1)(S^2 + 4)} \right]$

Consider

$$\frac{2S^2 - 1}{(S^2 + 1)(S^2 + 4)} = \frac{AS + B}{(S^2 + 1)} + \frac{CS + D}{(S^2 + 4)} \quad \text{--- (I)}$$

$$\frac{2S^2 - 1}{(S^2 + 1)(S^2 + 4)} = \frac{(AS + B)(S^2 + 4) + (CS + D)(S^2 + 1)}{(S^2 + 1)(S^2 + 4)}$$

$$2S^2 - 1 = (AS + B)(S^2 + 4) + (CS + D)(S^2 + 1)$$

$$2s^2 - 1 = As^3 + 4As + Bs^2 + 4B + Cs^2 + Cs + Ds^2 + D$$

$$2s^2 - 1 = s^3(A+C) + s^2(B+D) + s(4A+C) + 4B+D$$

Equating powers of s .

$$A+C=0, \quad B+D=2, \quad 4A+C=0, \quad 4B+D=-1$$

$$\therefore \boxed{A=0}$$

$$B+D=2$$

$$\boxed{C=0}$$

$$\begin{array}{r} 4B+D=-1 \\ - \quad - \quad + \end{array}$$

$$-3B = 3$$

$$\boxed{B=-1}$$

$$\therefore B+D=2$$

$$B(-1)+D=2$$

$$\boxed{D=3}$$

eqⁿ ② becomes

$$\frac{2s^2-1}{(s^2+1)(s^2+4)} = \frac{0 \times s + (-1)}{(s^2+1)} + \frac{0 \times s + 3}{(s^2+4)}$$

$$\frac{2s^2-1}{(s^2+1)(s^2+4)} = \frac{-1}{s^2+1} + \frac{3}{s^2+4}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{2s^2-1}{(s^2+1)(s^2+4)} \right] = L^{-1} \left[\frac{-1}{s^2+1} \right] + L^{-1} \left[\frac{3}{s^2+4} \right]$$

$$= -L^{-1}\left[\frac{1}{s^2+1}\right] + 3L^{-1}\left[\frac{1}{s^2+2^2}\right]$$

$$= -\frac{1}{1} \sin t + 3 \times \frac{1}{2} \sin 2t$$

$$\boxed{L^{-1}\left[\frac{2s^2-1}{(s^2+1)(s^2+4)}\right] = -\sin t + \frac{3}{2} \sin 2t}$$

3. Find $L^{-1}\left[\frac{1}{(s+1)(s^2+2s+2)}\right]$

Solⁿ:

Consider

$$\frac{1}{(s+1)(s^2+2s+2)} = \frac{A}{(s+1)} + \frac{Bs+C}{(s^2+2s+2)} \quad \text{--- (I)}$$

$$\frac{1}{(s+1)(s^2+2s+2)} = \frac{A(s^2+2s+2) + (Bs+C)(s+1)}{(s+1)(s^2+2s+2)}$$

$$1 = As^2 + 2As + 2A + Bs^2 + Bs + Cs + C$$

$$1 = s^2(A+B) + s(2A+B+C) + 2A+C$$

Equating powers of s

$$A+B=0, \quad 2A+B+C=0, \quad 2A+C=1$$

$$B=-A, \quad 2A+(-A)+C=0$$

$$A+C=0$$

$$2A + C = 1$$

$$A + C = 0$$

$$- \quad - \quad -$$

$$A + 0 = 1$$

$$\boxed{A = 1}$$

$$A + C = 0$$

$$C = -A$$

$$\boxed{C = -1}$$

$$B = -A$$

$$\therefore \boxed{B = -1}$$

eqⁿ ② becomes

$$\frac{1}{(s+1)(s^2+2s+2)} = \frac{1}{s+1} + \frac{(-1)s + (-1)}{(s^2+2s+2)}$$

$$\frac{1}{(s+1)(s^2+2s+2)} = \frac{1}{(s+1)} + \frac{-s-1}{(s^2+2s+2)}$$

$$= \frac{1}{s+1} - \frac{s+1}{s^2+2s+2}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{1}{(s+1)(s^2+2s+2)} \right] = L^{-1} \left[\frac{1}{s+1} \right] - L^{-1} \left[\frac{s+1}{s^2+2s+1+1} \right]$$

$$= L^{-1} \left[\frac{1}{s+1} \right] - L^{-1} \left[\frac{s+1}{(s+1)^2 + 1} \right]$$

$$= e^{-t} - e^{-t} L^{-1} \left[\frac{s}{s^2+1} \right]$$

$$= e^{-t} - e^{-t} \cos t$$

$$\therefore \mathcal{L}^{-1} \left[\frac{1}{(s+1)(s^2+2s+2)} \right] = e^{-t} [1 - \cos t]$$

4. Find $\mathcal{L}^{-1} \left[\frac{s}{(s+1)^2 (s^2+1)} \right]$

Consider,

$$\frac{s}{(s+1)^2 (s^2+1)} = \frac{A}{(s+1)} + \frac{B}{(s+1)^2} + \frac{Cs+D}{(s^2+1)} \quad \text{--- (I)}$$

$$s = A(s+1)(s^2+1) + B(s^2+1) + (Cs+D)(s+1)^2$$

$$s = A(s^3 + s + s^2 + 1) + B(s^2 + 1) + (Cs+D)(s^2 + 2s + 1)$$

$$s = As^3 + As + As^2 + A + Bs^2 + B + Cs^3 + 2Cs^2 + Cs + Ds^2 + 2Ds + D$$

$$s = s^3(A+C) + s^2(A+B+2C+D) + s(A+C+2D) + A+B+D$$

Equating powers of 's'

$$A+C=0, \quad A+B+2C+D=0, \quad A+C+2D=1, \quad A+B+D=0$$

$$\therefore C=-A \quad A+B+D+2C=0 \quad \text{Put } C=-A \quad 0+B+\frac{1}{2}=0$$

$$\therefore \boxed{A=0} \quad 0+2C=0 \quad A-A+2D=1$$

$$\boxed{C=0}$$

$$2D=1$$

$$\boxed{D=\frac{1}{2}}$$

$$\boxed{B=-\frac{1}{2}}$$

$\therefore$ eqⁿ (I) becomes

$$\frac{s}{(s+1)^2 (s^2+1)} = 0 + \frac{-\frac{1}{2}}{(s+1)^2} + \frac{\frac{1}{2}}{s^2+1}$$

Taking L^{-1} on both sides

$$L^{-1} \left[\frac{s}{(s+1)^2(s^2+1)} \right] = -\frac{1}{2} L^{-1} \left[\frac{1}{(s+1)^2} \right] + \frac{1}{2} L^{-1} \left[\frac{1}{s^2+1} \right]$$

$$= -\frac{1}{2} e^{-t} L^{-1} \left[\frac{1}{s^2} \right] + \frac{1}{2} L^{-1} \left[\frac{1}{s^2+1} \right]$$

$$= -\frac{1}{2} t e^{-t} + \frac{1}{2} \sin t$$

$$\therefore L^{-1} \left[\frac{s}{(s+1)^2(s^2+1)} \right] = \frac{1}{2} [\sin t - t e^{-t}]$$

~~Imp~~ #

Convolution Theorem

$$\mathcal{L}^{-1} [\bar{F}_1(s) \bar{F}_2(s)] = \int_0^t F_1(u) F_2(t-u) du$$

Examples

1. Solve by Convolution theorem

$$\mathcal{L}^{-1} \left[\frac{s}{(s^2+1)(s^2+4)} \right]$$

→ Let $\bar{F}_1(s) = \frac{1}{s^2+1}$

$$\bar{F}_2(s) = \frac{s}{s^2+4}$$

$$\mathcal{L}^{-1} [\bar{F}_1(s)] = \mathcal{L}^{-1} \left[\frac{1}{s^2+1} \right]$$

$$F_1(t) = \sin t$$

$$\therefore \boxed{F_1(u) = \sin u}$$

$$\bar{F}_2(s) = \frac{s}{s^2+4}$$

$$\mathcal{L}^{-1} [\bar{F}_2(s)] = \mathcal{L}^{-1} \left[\frac{s}{s^2+4} \right]$$

$$F_2(t) = \cos 2t$$

$$\therefore F_2(t-u) = \cos 2(t-u)$$

$$F_2(t-u) = \cos(2t-2u)$$

by Convolution Theorem

$$L^{-1}[\bar{F}_1(s) \bar{F}_2(s)] = \int_0^t F_1(u) F_2(t-u) du$$

$$L^{-1}\left[\frac{s}{(s^2+1)(s^2+4)}\right] = \int_0^t \sin u \cdot \cos(2t-2u) du$$

$$= \frac{1}{2} \int_0^t 2 \sin u \cdot \cos(2t-2u) du$$

$$2 \sin A \cdot \cos B = \sin(A+B) + \sin(A-B)$$

$$= \frac{1}{2} \int_0^t [\sin(u+2t-2u) + \sin(u-(2t-2u))] du$$

$$= \frac{1}{2} \int_0^t \sin(2t-u) + \sin(3u-2t) du$$

$$= \frac{1}{2} \left[\frac{-\cos(2t-u)}{(0-1)} + \left(\frac{-\cos(3u-2t)}{(3-0)} \right) \right]_0^t$$

$$= \frac{1}{2} \left[\cos(2t-t) - \cos(3t-2t) - \right.$$

$$\left. \left(\frac{\cos(2t-0)}{3} - \cos(0-2t) \right) \right]$$

$$= \frac{1}{2} \left[\cos t - \frac{\cos t}{3} - \left[\cos 2t - \frac{\cos 2t}{3} \right] \right]$$

$$= \frac{1}{2} \left[\frac{3 \cos t - \cos t}{3} - \left[\frac{3 \cos 2t - \cos 2t}{3} \right] \right]$$

$$= \frac{1}{2} \left[\frac{2 \cos t}{3} - \frac{2 \cos 2t}{3} \right]$$

$$= \frac{2}{6} [\cos t - \cos 2t]$$

$$= \frac{1}{3} [\cos t - \cos 2t]$$

$$\therefore \mathcal{L}^{-1} \left[\frac{s}{(s^2+1)(s^2+4)} \right] = \frac{1}{3} [\cos t - \cos 2t]$$

2 By using convolution theorem Solve
 $\mathcal{L}^{-1} \left[\frac{1}{s(s+1)(s+2)} \right]$

→ Solⁿ, Let $\bar{F}_1(s) = \frac{1}{s(s+1)}$

$$\bar{F}_2(s) = \frac{1}{s+2}$$

$$\mathcal{L}^{-1} [\bar{F}_1(s)] = \mathcal{L}^{-1} \left[\frac{1}{s(s+1)} \right]$$

$$F_1(t) = \int_0^t e^{-t} dt = \left[\frac{e^{-t}}{-1} \right]_0^t$$

$$F_1(t) = 1 - e^{-t}$$

$$F_1(u) = 1 - e^{-u}$$

$$\bar{F}_2(s) = \frac{1}{s+2}$$

$$L^{-1}[\bar{F}_2(s)] = L^{-1}\left[\frac{1}{s+2}\right]$$

$$F_2(t) = e^{-2t}$$

$$F_2(t-u) = e^{-2(t-u)}$$

$$\therefore F_2(t-u) = e^{-2t+2u}$$

by convolution Th^m

$$L^{-1}[\bar{F}_1(s) \bar{F}_2(s)] = \int_0^t F_1(u) \cdot F_2(t-u) du$$

$$L^{-1}\left[\frac{1}{s(s+1)(s+2)}\right] = \int_0^t (1-e^{-u}) e^{-2t+2u} du$$

$$= \int_0^t (1-e^{-u}) e^{-2t} e^{2u} du$$

$$= e^{-2t} \int_0^t (1-e^{-u}) e^{2u} du$$

$$= e^{-2t} \int_0^t (e^{2u} - e^{2u} e^{-u}) du$$

$$= e^{-2t} \int_0^t (e^{2u} - e^u) du$$

$$= e^{-2t} \left[\frac{e^{2t}}{2} - e^t \right]_0^t$$

$$= e^{-2t} \left[\frac{e^{2t}}{2} - e^t - \left(\frac{e^0}{2} - e^0 \right) \right]$$

$$= e^{-2t} \left[\frac{e^{2t}}{2} - e^t - \left(\frac{1}{2} - 1 \right) \right]$$

$$= e^{-2t} \left[\frac{e^{2t}}{2} - e^t - \left(-\frac{1}{2} \right) \right]$$

$$= e^{-2t} \left[\frac{e^{2t}}{2} - e^t + \frac{1}{2} \right]$$

Solution of different
H.W.

$$3. \quad L^{-1} \left[\frac{s^2}{(s^2+a^2)(s^2+b^2)} \right]$$

$$\rightarrow \text{Let } \bar{F}_1(s) = \frac{s}{s^2+a^2}, \quad \bar{F}_2(s) = \frac{s}{s^2+b^2}$$

$$L^{-1}[\bar{F}_1(s)] = L^{-1} \left[\frac{s}{s^2+a^2} \right], \quad L^{-1}[\bar{F}_2(s)] = L^{-1} \left[\frac{s}{s^2+b^2} \right]$$

$$F_1(t) = \cos at$$

$$F_2(t) = \cos bt$$

$$\therefore \boxed{F_1(u) = \cos au}$$

$$\therefore \boxed{F_2(t-u) = \cos(bt-bu)}$$

by Convolution theorem,

$$L^{-1}[\bar{F}_1(s) \cdot \bar{F}_2(s)] = \int_0^t F_1(u) \cdot F_2(t-u) du$$

$$\therefore \mathcal{L}^{-1} \left[\frac{s^2}{(s^2+a^2)(s^2+b^2)} \right] = \int_0^t \cos au \cdot \cos(bt-bu) du$$

$$= \frac{1}{2} \int_0^t 2 \cos au \cdot \cos(bt-bu) du$$

$$= \frac{1}{2} \int_0^t [\cos(au+bt-bu) + \cos(au-bt+bu)] du$$

$$= \frac{1}{2} \left[\frac{\sin((a-b)u+bt)}{a-b} + \frac{\sin((a+b)u-bt)}{a+b} \right]_0^t$$

$$= \frac{1}{2} \left[\frac{\sin((a-b)t+bt)}{a-b} + \frac{\sin((a+b)t-bt)}{a+b} \right]$$

$$- \left(\frac{\sin((a-b)0+bt)}{a-b} + \frac{\sin((a+b)0-bt)}{a+b} \right)$$

$$= \frac{1}{2} \left[\frac{\sin(at-bt+bt)}{a-b} + \frac{\sin(at+bt-bt)}{a+b} \right]$$

$$- \left(\frac{\sin(0+bt)}{a-b} + \frac{\sin(0-bt)}{a+b} \right)$$

$$= \frac{1}{2} \left[\frac{\sin at}{a-b} + \frac{\sin at}{a+b} - \left(\frac{\sin bt}{a-b} - \frac{\sin bt}{a+b} \right) \right]$$

$$= \frac{1}{2} \left[\sin at \left(\frac{1}{a-b} + \frac{1}{a+b} \right) - \sin bt \left(\frac{1}{a-b} - \frac{1}{a+b} \right) \right]$$

$$= \frac{1}{2} \left[\sin at \left(\frac{2a}{a^2-b^2} \right) - \sin bt \left(\frac{2b}{a^2-b^2} \right) \right]$$

$$\therefore \mathcal{L}^{-1} \left[\frac{s^2}{(s^2+a^2)(s^2+b^2)} \right] = \frac{a \sin at - b \sin bt}{a^2-b^2}$$

95%
#

Solution of differential equation with Constant Coefficient.

$$L[F'(t)] = s\bar{F}(s) - F(0)$$

$$L[F''(t)] = s^2\bar{F}(s) - sF(0) - F'(0)$$

$$L[F'''(t)] = s^3\bar{F}(s) - s^2F(0) - sF'(0) - F''(0)$$

Examples

1. $\frac{d^3y}{dt^3} + 2\frac{d^2y}{dt^2} - \frac{dy}{dt} - 2y = 0$

$$y = 1, \frac{dy}{dt} = 2, \frac{d^2y}{dt^2} = 2 \text{ at } t = 0$$

Solⁿ. We have

$$\frac{d^3y}{dt^3} + 2\frac{d^2y}{dt^2} - \frac{dy}{dt} - 2y = 0$$

$$y'''(t) + 2y''(t) - y'(t) - 2y(t) = 0 \quad \text{--- (A)}$$

$$\text{at } t = 0, y(0) = 1, y'(0) = 2, y''(0) = 2$$

Taking L.T on both eqⁿ (A)

$$L[y'''(t)] + 2L[y''(t)] - L[y'(t)] - 2L[y(t)] = 0$$

$$s^3 \bar{y}(s) - s^2 y(0) - s y'(0) - y''(0) + 2[s^2 \bar{y}(s) - s y(0) - y'(0)] - [s \bar{y}(s) - y(0)] - 2 \bar{y}(s) = 0$$

$$s^3 \bar{y}(s) - s^2(1) - s(2) - 2 + 2s^2 \bar{y}(s) - 2s(1) - 2(2) - s \bar{y}(s) + 1 - 2 \bar{y}(s) = 0$$

$$\bar{y}(s) [s^3 + 2s^2 - s - 2] - s^2 - 2s - 2 - 2s - 4 + 1 = 0$$

$$\bar{y}(s) (s^3 + 2s^2 - s - 2) - s^2 - 4s - 5 = 0$$

$$\bar{y}(s) [s^3 + 2s^2 - s - 2] = s^2 + 4s + 5$$

$$\bar{y}(s) = \frac{s^2 + 4s + 5}{s^3 + 2s^2 - s - 2} \rightarrow \textcircled{B}$$

Consider,

$$s^3 + 2s^2 - s - 2 \quad \begin{array}{c|cccc} 1 & 1 & 2 & -1 & -2 \\ & & 1 & 1 & -2 \\ \hline & 1 & 1 & -2 & 0 \end{array}$$

$$(s+1)(s^2 + s - 2) = 0$$

$$(s+1)(s+2)(s-1) = 0$$

eqⁿ $\textcircled{B}$ becomes

$$\bar{y}(s) = \frac{s^2 + 4s + 5}{s^3 + 2s^2 - s - 2} = \frac{s^2 + 4s + 5}{(s+1)(s-1)(s+2)}$$

$$= \frac{A}{(s+1)} + \frac{B}{(s-1)} + \frac{C}{(s+2)}$$

(c)

$$s^2 + 4s + 5 = A(s-1)(s+2) + B(s+1)(s+2) + C(s-1)(s+1)$$

$$= A(s^2 + 2s - s - 2) + B(s^2 + 2s + s + 2) + C(s^2 + s - s - 1)$$

$$= A(s^2 + s - 2) + B(s^2 + 3s + 2) + C(s^2 - 1)$$

$$= As^2 + As - 2A + Bs^2 + 3Bs + 2B + Cs^2 - C$$

$$s^2 + 4s + 5 = s^2(A+B+C) + s(A+3B) + 2B - C - 2A$$

$$\therefore A+B+C=1, \quad A+3B=4, \quad -2A+2B-C=5$$

$$-2A+2B-C=5$$

$$A+B+C=1$$

$$-A+3B+0=6$$

$$A+3B=4$$

$$-A+3B=6$$

$$6B=10$$

$$\boxed{B = \frac{5}{3}}$$

$$A+3B=4$$

$$A=4-3B$$

$$= 4 - 3 \times \frac{5}{3}$$

$$= 4 - 5$$

$$\boxed{A = -1}$$

$$\therefore A+B+C=1$$

$$-1 + \frac{5}{3} + C = 1$$

$$\therefore C = 1 + 1 - \frac{5}{3}$$

$$C = \frac{2-5}{3}$$

$$C = \frac{6-5}{3}$$

$$C = \frac{1}{3}$$

eqⁿ (c) becomes

$$\frac{s^2 + 4s + 5}{s^3 + 2s^2 - s - 2} = \frac{-1}{(s+1)} + \frac{5}{3} \times \frac{1}{s-1} + \frac{1}{3} \times \frac{1}{s+2}$$

L^{-1} on both sides

$$L^{-1} [\gamma(s)] = L^{-1} \left[\frac{s^2 + 2s + 5}{s^3 + 2s^2 - s - 2} \right] = -1 L^{-1} \left[\frac{1}{s+1} \right] + \frac{5}{3} L^{-1} \left[\frac{1}{s-1} \right] + \frac{1}{3} L^{-1} \left[\frac{1}{s+2} \right]$$

$$\gamma(t) = -e^{-t} + \frac{5}{3} e^t + \frac{1}{3} e^{-2t}$$

2. $\frac{d^2 x}{dt^2} + 9x = \cos 2t$ if $x(0) = 1$, $x\left(-\frac{\pi}{2}\right) = -1$


→ We have,

$$\frac{d^2 x}{dt^2} + 9x = \cos 2t$$

$$x''(t) + 9x(t) = \cos 2t \text{ if } x(0) = 1,$$

Taking L.T on both sides

$$\frac{-1+5}{5} = \frac{4}{5}$$



$$L[x''(t)] + gL[x(t)] = L[\cos 2t]$$

$$s^2 \bar{x}(s) - s(0) - x'(0) + g\bar{x}(s) = \frac{s}{s^2+4}$$

$$s^2(\bar{x}(s)) - s(1) - k + g\bar{x}(s) = \frac{s}{s^2+4}$$

$$\bar{x}(s)(s^2+g) - s - k = \frac{s}{s^2+4}$$

$$\bar{x}(s)(s^2+g) = \frac{s}{s^2+4} + s + k$$

$$\bar{x}(s) = \frac{s}{(s^2+4)(s^2+g)} + \frac{s}{s^2+g} + \frac{k}{s^2+g}$$

Taking L^{-1} on both sides

$$L^{-1}[\bar{x}(s)] = L^{-1}\left[\frac{s}{(s^2+4)(s^2+g)}\right] + L^{-1}\left[\frac{s}{s^2+g}\right] + L^{-1}\left[\frac{k}{s^2+g}\right]$$

$$= L^{-1}\left[\frac{s}{(s^2+4)(s^2+g)}\right] + \cos 3t + \frac{k \sin 3t}{3}$$

A_1

— (A)

Consider

$$A_1 = \frac{s}{(s^2+4)(s^2+g)} = \frac{As+B}{(s^2+4)} + \frac{Cs+D}{(s^2+g)} \quad \text{--- (B)}$$

$$\frac{s}{(s^2+4)(s^2+g)} = \frac{(As+B)(s^2+g) + (Cs+D)(s^2+4)}{(s^2+4)(s^2+g)}$$

$$S = AS^3 + 9As + BS^2 + 9B + CS^3 + 4Cs + DS^2 + 4D$$

$$S = S^3[A+C] + S^2[B+D] + S[9A+4C] + 9B+4D$$

Equating powers of s

$$A+C=0, B+D=0, 9A+4C=1, 9B+4D=0$$

$$A=-C$$

$$B=-D$$

$$\text{Put } A=-C$$

$$9(-D)+4D=0$$

$$\text{Put } C=-\frac{1}{5}$$

$$\text{Put } D=0$$

$$9(-C)+4C=1$$

$$-9D+4D=0$$

$$\therefore \boxed{B=0}$$

$$-9C+4C=1$$

$$-5D=0$$

$$-5C=1$$

$$\therefore \boxed{D=0}$$

$$\therefore \boxed{A = \frac{1}{5}}$$

$$\boxed{C = -\frac{1}{5}}$$

eqⁿ (B) becomes

$$A_1 = \frac{S}{(S^2+4)(S^2+9)} = \frac{\frac{1}{5}S+0}{S^2+4} + \frac{-\frac{1}{5}S+0}{S^2+9}$$

$$= \frac{1}{5} \frac{S}{S^2+4} - \frac{1}{5} \frac{S}{S^2+9}$$

$$= \frac{1}{5} \mathcal{L}^{-1} \left[\frac{S}{S^2+2^2} \right] - \frac{1}{5} \mathcal{L}^{-1} \left[\frac{S}{S^2+3^2} \right]$$

$$= \frac{1}{5} \cos 2t - \frac{1}{5} \cos 3t$$

eqⁿ (A) becomes

$$\mathcal{L}^{-1}[\bar{x}(s)] = \frac{1}{5} \cos 2t - \frac{1}{5} \cos 3t + \cos 3t + \frac{k}{3} \sin 3t$$

(A)

$$x(t) = \frac{1}{5} \cos 2t + \frac{4 \cos 3t}{5} + \frac{k}{3} \sin 3t \quad \text{--- (c)}$$

$$x\left(\frac{\pi}{2}\right) = -1 \quad \text{put } t = \frac{\pi}{2}$$

$$x\left(\frac{\pi}{2}\right) = \frac{1}{5} \cos 2 \times \frac{\pi}{2} + \frac{4 \cos\left(\frac{3\pi}{2}\right)}{5} + \frac{k}{3} \sin\left(\frac{3\pi}{2}\right)$$

$$-1 = \frac{1}{5} (-1) + \frac{4 \times 0}{5} + \frac{k}{3} (-1)$$

$$-1 = -\frac{1}{5} - \frac{k}{3}$$

$$\frac{k}{3} = -\frac{1}{5} + 1$$

$$\boxed{\frac{k}{3} = \frac{4}{5}} \quad \therefore \boxed{k = \frac{12}{5}}$$

eqⁿ (c) becomes

$$\boxed{x(t) = \frac{1}{5} \cos 2t + \frac{4 \cos 3t}{5} + \frac{4 \sin 3t}{5}}$$

Solution of Simultaneous L.D.E

* Examples

1. Solve by L.T $\frac{dx}{dt} - y = e^t$, $\frac{dy}{dt} + x = \sin t$

given that $x(0) = 1$, $y(0) = 0$

⇒ Solⁿ, we have

$$\frac{dx}{dt} - y = e^t$$

$$x'(t) - y(t) = e^t \quad \text{--- (i)}$$

$$L[x'(t)] - L[y(t)] = L[e^t]$$

$$s\bar{x}(s) - x(0) - \bar{y}(s) = \frac{1}{s-1}$$

$$s\bar{x}(s) - \bar{y}(s) - 1 = \frac{1}{s-1}$$

$$s\bar{x}(s) - \bar{y}(s) = \frac{1}{s-1} + 1$$

$$= \frac{1 + s - 1}{s-1}$$

$$s\bar{x}(s) - \bar{y}(s) = \frac{s}{s-1} \rightarrow \textcircled{A}$$

Again consider

$$\frac{dy}{dt} + x = \sin t$$

$$y'(t) + x(t) = \sin t$$

$$L[y'(t)] + L[x(t)] = L[\sin t]$$

$$s\bar{y}(s) + y(0) + \bar{x}(s) = \frac{1}{s^2+1}$$

$$s\bar{y}(s) + \bar{x}(s) - 0 = \frac{1}{s^2+1}$$

$$s\bar{y}(s) + \bar{x}(s) = \frac{1}{s^2+1}$$

$$\bar{x}(s) + s\bar{y}(s) = \frac{1}{s^2+1} \quad \text{--- (B)}$$

multiply by 's' to eqⁿ A

$$s^2\bar{x}(s) - s\bar{y}(s) = \frac{s^2}{s^2+1}$$

$$+ \bar{x}(s) + s\bar{y}(s) = \frac{1}{s^2+1}$$

$$s^2\bar{x}(s) + \bar{x}(s) = \frac{s^2}{s^2+1} + \frac{1}{s^2+1}$$

$$\bar{x}(s)(s^2+1) = \frac{s^2}{s^2+1} + \frac{1}{s^2+1}$$

$$\mathcal{L}^{-1}[\bar{x}(s)] = \mathcal{L}^{-1}\left[\frac{s^2}{(s-1)(s^2+1)}\right] + \mathcal{L}^{-1}\left[\frac{1}{(s^2+1)(s^2+1)}\right] \quad \text{--- (c)}$$

C_1 C_2

Consider,

$$C_1 = \frac{s^2}{(s-1)(s^2+1)} = \frac{A}{s-1} + \frac{Bs+C}{s^2+1}$$

$$s^2 = A(s^2+1) + (Bs+C)(s-1)$$

$$= As^2 + A + Bs^2 - Bs + Cs - C$$

$$s^2 = s^2(A+B) + s(-B+C) + A-C$$

Equating powers of s

$$A+B=1, \quad -B+C=0, \quad A-C=0$$

$$C=B, \quad A-B=0$$

$$A+B=1$$

$$A-B=0$$

$$2A=1$$

$$\therefore \boxed{A = \frac{1}{2}}$$

$$\therefore \boxed{B = \frac{1}{2}}$$

$$\therefore \boxed{C = \frac{1}{2}}$$

$$\therefore \mathcal{L}^{-1}\left[\frac{s^2}{(s-1)(s^2+1)}\right] = \mathcal{L}^{-1}\left[\frac{1}{2} \frac{1}{s-1} + \frac{1}{2} \frac{s+1}{s^2+1}\right]$$

$$= \frac{1}{2} \mathcal{L}^{-1} \left[\frac{1}{s-1} \right] + \frac{1}{2} \mathcal{L}^{-1} \left[\frac{s}{s^2+1} \right] + \frac{1}{2} \mathcal{L}^{-1} \left[\frac{1}{s^2+1} \right]$$

$$\boxed{\mathcal{L}^{-1} \left[\frac{s^2}{(s-1)(s^2+1)} \right] = \frac{1}{2} e^t + \frac{1}{2} \cos t + \frac{1}{2} \sin t} \quad \text{--- } C_1$$

Consider,

$$C_2 = \frac{1}{(s^2+1)(s^2+1)} = \frac{As+B}{(s^2+1)} + \frac{Cs+D}{(s^2+1)^2}$$

$$\frac{1}{(s^2+1)^2} = \frac{(s^2+1)(As+B) + (Cs+D)}{(s^2+1)^2}$$

$$1 = (s^2+1)(As+B) + (Cs+D)$$

$$1 = As^3 + Bs^2 + As + B + Cs + D$$

$$1 = As^3 + Bs^2 + s(CA+C) + B+D$$

Equating powers of s

$$\boxed{A=0}, \boxed{B=0}, A+C=0, B+D=1$$

$$0+C=0$$

$$0+D=1$$

$$\therefore \boxed{C=0}$$

$$\therefore \boxed{D=1}$$

$$\therefore C_2 = \frac{1}{(s^2+1)^2} = \frac{0+0}{(s^2+1)} + \frac{1}{(s^2+1)^2}$$

$$\mathcal{L}^{-1} \left[\frac{1}{(s^2+1)^2} \right]$$

Consider,

$$C_2 = \frac{1}{(s^2+1)^2}$$

$$\text{let } \bar{F}_1(s) = \frac{1}{s^2+1}, \quad \bar{F}_2(s) = \frac{1}{s^2+1}$$

$$\mathcal{L}^{-1}[\bar{F}_1(s)] = \sin t, \quad \mathcal{L}^{-1}[\bar{F}_2(s)] = \sin t$$
$$F_1(t) = \sin t, \quad F_2(t) = \sin t$$

$$\therefore \boxed{F_1(u) = \sin u}, \quad \boxed{F_2(u) = \sin(t-u)}$$

by

By convolution theorem

$$\mathcal{L}^{-1}[\bar{F}_1(s) \cdot \bar{F}_2(s)] = \int_0^t F_1(u) \cdot F_2(t-u) du$$
$$= \int_0^t \sin u \sin(t-u) du$$

use $2\sin A \cdot \sin B = \cos(A-B) - \cos(A+B)$

$$= \frac{1}{2} \int_0^t 2 \sin u \sin(t-u) du$$

$$= \frac{1}{2} \int_0^t (\cos(u-(t-u)) - \cos(u+(t-u))) du$$

$$= \frac{1}{2} \int_0^t (\cos(2u-t) - \cos(t)) du$$

$$= \frac{1}{2} \left[\frac{\sin(2u-t)}{2} - \cos t [u] \right]_0^t$$

$$= \frac{1}{2} \left[\frac{\sin t - t \cos t}{2} - \left(\frac{\sin(-t)}{2} - 0 \right) \right]$$

$$= \frac{1}{2} \left[\frac{\sin t - t \cos t}{2} - \left(\frac{-\sin t}{2} \right) \right]$$

$$= \frac{1}{2} \left[\frac{\sin t + \sin t - t \cos t}{2} \right]$$

$$= \frac{1}{2} \left[\frac{2 \sin t - t \cos t}{2} \right]$$

$$\mathcal{L}^{-1} \left[\frac{1}{(s^2+1)^2} \right] = \frac{1}{2} \sin t - \frac{t}{2} \cos t \quad \text{--- } C_2$$

$\therefore$ eqⁿ (C) becomes.

$$\boxed{x(t) = \frac{1}{2} e^t + \frac{1}{2} \cos t + \sin t - \frac{t}{2} \cos t} \quad \text{--- (D)}$$

Diff. w.r.t. 't'

$$\therefore x'(t) = \frac{1}{2} e^t - \frac{1}{2} \sin t + \cos t - \frac{1}{2} [-t \sin t + \cos t]$$

$$\boxed{x'(t) = \frac{1}{2} e^t - \frac{1}{2} \sin t + \frac{1}{2} \cos t + \frac{t}{2} \sin t} \quad \text{--- (2)}$$

put eqⁿ (2) in eqⁿ (1) we get

$$\boxed{y(t) = -\frac{1}{2} e^t - \frac{1}{2} \sin t + \frac{1}{2} \cos t + \frac{t}{2} \sin t}$$